

| DIRECT MEASURING BY WEIGHT | PRO PROGRAM | 3-Part System (Micro, Grow, Bloom)

< PRO > FEED PROGRAM							In gram(g.) per 1 Liter of water MEASURING BY WEIGHT									
Mixing Order		Clone		Vegetative			Flower									
		First 7 Days	Day 8 to plant	W1	W2	WЗ	W4	W1	W2	WЗ	W4	W5	W6	W7	W8	Flush
← Mixing Order	CoreX (Micro, Core)	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	
	EldeX (Grow)	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	
	FloreX (Bloom)	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.25
	Aurora (PK Amino)									0.15	0.15	0.25	0.25	0.25	0.25	
	Mira (pH UP Silica)	Put a few drops to boost up the pH (If the pH is optimal without Mira then you don't need)														
Targets	EC	1.8	1.8	1.8	1.8	1.8	1.8	1.8	1.8	2.1	2.1	2.4	2.4	2.4	2.2	0.1
	PPM 500	900	900	900	900	900	900	900	900	1050	1050	1200	1200	1200	1100	50
	PPM 700	1260	1260	1260	1260	1260	1260	1260	1260	1470	1470	1680	1680	1680	1540	70
	рН	5.6	5.8	5.5-5.8 (Coco/Rockwool) 5.9-6.2 (Peat moss base)				5.8-6.2 (Coco/Rockwool) 6.0-6.4 (Peat moss base)							6.0 - 6.4	

- 1. Runoff every time.
- 2. If strains go past Week 8, continue same schedule as Week 8.
- 3. Flush before harvest.

The target EC (or PPM) can differ by about 10%.

Runoff

Veg, Flower 5-10% runoff, dry back 50% of water on first run and 35% dry back all the time

Flush 30%+ runoff, dry back 15% of water

3-Part ENG



SCAN FOR MORE DETAIL

SPRA		Vege	tative	Flower				
Applica	W1	W2	wз	W4	W1	W2	МЗ	
Terra (Humic,Kelp)	2X Week	0.25g	0.25g	0.25g	0.25g	0.25g	0.25g	0.25g

All measurements are Gram (g) per Liter (l)

PRE-SOAK

Coco
pH 5.5-5.8
EC 3.5-4.0

Rockwool
pH 5.0-5.5
EC 2.5-3.0

Rockwool
pH 5.0-5.5

FLUSH
pH 6.0-6.4
Peat moss
Last 5 days
Coco
Last 3 days
Rockwool
Last days

Note: The feed chart should be adjusted based on the duration of the run, which can vary depending on the strain being used.