

# FOR STOCK TANK STARTER PROGRAM 3-Part System (Micro, Grow, Bloom)

< STARTER > FEED PROGRAM								Based on 250 grams in 1 Liter stock tank concentrate. All measurements are milliliters (mL) per Liter (l)								
Mixing Order		Clone		Vegetative				Flower								
		First 7 Days	Day 8 to plant	W1	W2	W3	W4	W1	W2	W3	W4	W5	W6	W7	W8	Flush
Mixing Order	CoreX (Micro, Core)	3.0	3.0	2.0	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	
	EldeX (Grow)	3.0	3.0	2.0	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	
	FloreX (Bloom)	3.0	3.0	2.0	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5	1.0
	Aurora (PK Amino)									0.6	0.6	0.6	0.6	0.6	0.6	
	Mira (pH UP Silica)	Put a few drops to boost up the pH (If the pH is optimal without Mira then you don't need)														
Targets	EC	1.8	1.8	1.3	1.5	1.5	1.5	1.5	1.5	1.9	1.9	1.9	1.9	1.9	1.9	0.1
	PPM 500	900	900	650	750	750	750	750	750	950	950	950	950	950	950	50
	PPM 700	1260	1260	910	1050	1050	1050	1050	1050	1330	1330	1330	1330	1330	1330	70
	pH	5.6	5.8	5.5-5.8 (Coco/Rockwool) 5.9-6.2 (Peat moss base)				5.8-6.2 (Coco/Rockwool) 6.0-6.4 (Peat moss base)								6.0 - 6.4

1. Runoff every time.
2. If strains go past Week 8, continue same schedule as Week 8.
3. Flush before harvest.

The target EC (or PPM) can differ by about 10%.

3-Part ENG



SCAN FOR MORE DETAIL

Runoff	
Veg, Flower	5-10% runoff, dry back 50% of water on first run and 35% dry back all the time
Flush	30%+ runoff, dry back 15% of water

SPRAY PROGRAM		Vegetative				Flower		
Application Frequency		W1	W2	W3	W4	W1	W2	W3
Terra (Humic, Kelp)	2X Week	0.25g	0.25g	0.25g	0.25g	0.25g	0.25g	0.25g

All measurements are Gram (g) per Liter (l)

PRE-SOAK
Coco pH 5.5-5.8 EC 3.5-4.0
Rockwool pH 5.0-5.5 EC 2.5-3.0

FLUSH
pH 6.0-6.4
Peat moss Last 5 days
Coco Last 3 days
Rockwool Last days

Note: The feed chart should be adjusted based on the duration of the run, which can vary depending on the strain being used.