

Concentrated Liquid Feed - 2 Parts (AB)

2 Parts F	eeding Chart	Baseline ratio 1:2							
	Shorten the mixing process by separating the Vegetative state (COREX + ELDEX) and the Flowering state (COREX + FLOREX), using only 2 parts (A and B)								
Concen	Concentrated Liquid Feed								
		'	od (b.1) for measuring by volume with a measuring cup uitable for automatic fertilizer dispensers.						
Making concentrated liquid fertilizer									
	•	J .	liquid fertilizer as per (b.1) to convert powder/granule/according to this feeding chart.						

Feeding Chart XC2		KING WHALE- HIGH EC						
		Vegetative state			Flowering state			
Mixing Order	Units	Clone	First week	Every week	Week 1 - 2	Week 3 - 6	Week 7 - 8+	Flush
COREX - Baseline Part A	ml./L	2.8	4	4	4	4		
ELDEX - Baseline Part B	ml./L	2.8	8	8				
FLOREX - Baseline Part B	ml./L	2.8			8	8	8	
FLO FADE - Late Color Booster	ml./L						4	2
AURORA - Flower Booster	ml./L				2	4	2	
TERRA - Root & Bud Formation	g./L	0.1						
OXYROOT - Less Buildup	ml./L	0.5 - 1 ml. All stage						
MIRA UP - Stalk & Leaf Thicker	ml./L	0.1 - 1 ml. All Stage, Use to increase pH, check pH value as need						
EC		1.8	2.8	2.8	3.2	3.7	3.5	0.3
PPM 500		900	1400	1400	1600	1850	1750	150
PPM 700		1260	1960	1960	2240	3330	3150	210

Foliar spray				
. Gilai Spiaj	Vegetative state			
Mixing Order	Units	First week	Every week	
TERRA - Root & Bud Formation	g./L	0.1	0.1	

Check pH and EC every time

To ensure accurate fertilizer mixing Optimal pH range is between 5.5 - 6.5

pH too low::

- 1. Reduce concentration to match water used
- 2. Use **MIRA UP** to increase pH

pH too high:

1. Increase fertilizer concentration

