

Concentrated Liquid Feed - 3 Parts

3 Parts Feeding Chart Baseline ratio 1:1:1 Use baseline fertilizers (*COREX* + ELDEX* + FLOREX**) during all stages without changing fertilizers, and use them in equal proportions. This simplifies fertilizer stock management and reduces mixing confusion. Concentrated Liquid Feed Use concentrated liquid fertilizer from method (b.1) for measuring by volume with a measuring cup (no need to weigh). Convenient for use and suitable for automatic fertilizer dispensers. Making concentrated liquid fertilizer Study the method for making concentrated liquid fertilizer as per (b.1) to convert powder/granule/

flakes into concentrated liquid form for use according to this feeding chart.

Feeding Chart XC3		KING WHALE- HIGH EC							
		Vegetative state			Flowering state				
Mixing Order	Units	Clone	First week	Every week	Week 1 - 2	Week 3 - 6	Week 7 - 8+	Flush	
COREX - Baseline Part A	ml./L	2.8	4	4	4	4			
ELDEX - Baseline Part B	ml./L	2.8	4	4	4	4	4		
FLOREX - Baseline Part B	ml./L	2.8	4	4	4	4	4		
FLO FADE - Late Color Booster	ml./L						4	2	
AURORA - Flower Booster	ml./L				2	4	2		
TERRA - Root & Bud Formation	g./L	0.1							
OXYROOT - Less Buildup	ml./L	0.5 - 1 ml. All stage							
MIRA UP - Stalk & Leaf Thicker	ml./L	0.1 - 1 ml. All Stage, Use to increase pH, check pH value as need							
EC		1.8	2.8	2.8	3.2	3.7	3.5	0.3	
PPM 500		900	1400	1400	1600	1850	1750	150	
PPM 700		1260	1960	1960	2240	3330	3150	210	

Foliar spray	Vegetative state			
Mixing Order	Units	First week	Every week	
TERRA - Root & Bud Formation	g./L	0.1	0.1	

Check pH and EC every time

To ensure accurate fertilizer mixing Optimal pH range is between 5.5 - 6.5

nH too low.

- Reduce concentration to match water used
- 2. Use $\emph{MIRA UP}$ to increase pH

pH too high:

1. Increase fertilizer concentration

